

Supplementary data for the article:

Ristivojević, P.; Dimkić, I.; Guzelmeric, E.; Trifković, J.; Knežević, M.; Berić, T.; Yesilada, E.; Milojković-Opsenica, D.; Stanković, S. Profiling of Turkish Propolis Subtypes: Comparative Evaluation of Their Phytochemical Compositions, Antioxidant and Antimicrobial Activities. *LWT* **2018**, *95*, 367–379.
<https://doi.org/10.1016/j.lwt.2018.04.063>

Supplementary material

Profiling of Turkish propolis subtypes: comparative evaluation of their phytochemical compositions, antioxidant and antimicrobial activities

Petar Ristivojević^{1,2¶*}, Ivica Dimkić^{3¶*}, Etil Guzelmeric⁴, Jelena Trifkovic⁵, Magdalena Knežević⁶, Tanja Berić³, Erdem Yesilada⁴, Dušanka Milojković-Opsenica⁵, Slaviša Stanković³

¹University of Belgrade - Innovation Centre of the Faculty of Chemistry Ltd., Studentski trg 12-16, 11000 Belgrade, Serbia

²Ghent University Global Campus, Incheon, South Korea

³University of Belgrade - Faculty of Biology, Studentski trg 16, 11000 Belgrade, Serbia

⁴Yeditepe University - Faculty of Pharmacy, Department of Pharmacognosy and Phytotherapy,

Kayisdagi Cad., Atasehir, 34755, Istanbul, Turkey

⁵University of Belgrade - Faculty of Chemistry, P.O. Box 51, 11158 Belgrade, Serbia

⁶Institute of Soil Science, Teodora Drajzera 7, 11000 Belgrade, Serbia

¶These authors contributed equally to this work.

*Corresponding author:

Ivica Dimkić, PhD

Research associate

University of Belgrade - Faculty of Biology,

Studentski trg 16, 11000 Belgrade, Serbia

E-mail: ivicad@bio.bg.ac.rs

Petar Ristivojević, PhD

Research associate

University of Belgrade - Innovation Centre of the Faculty of Chemistry Ltd.,

Studentski trg 12-16, 11000 Belgrade, Serbia

E-mail: ristivojevic@chem.bg.ac.rs

List of figures

Fig. S1. Propolis samples collected from different regions of Turkey

Fig. S2. RSA values of Turkish propolis samples

Fig. S3. The calibration growth curves for: *C. albicans* (A), *S. mutans* (B), *S. pyogenes* (C) and *S. sanguinis* (D). On the right ordinate, optical density values measured at time interval of 8 hours up to the monitored time are displayed. The left ordinate represents the logarithmic values of CFU/mL.



Fig. S1. Propolis samples collected from different regions of Turkey




Fig. S2. RSA values of Turkish propolis samples




Fig. S3. The calibration growth curves for: *C. albicans* (A), *S. mutans* (B), *S. pyogenes* (C) and *S. sanguinis* (D). On the right ordinate, optical density values measured at time interval of 8 hours up to the monitored time are displayed. The left ordinate represents the logarithmic values of CFU/mL.